



Corporate Presentation

April 2023



Safe Harbor Statement

This presentation includes forward-looking statements about, among other things, Verastem Oncology's programs and product candidates, including anticipated regulatory submissions, approvals, performance and potential benefits of Verastem Oncology's product candidates, as well as Verastem Oncology's potential income under its asset purchase agreement with Secura Bio. Inc. and borrowings under its credit facility, that are subject to substantial risks and uncertainties that could cause actual results to differ materially from those expressed or implied by such statements. Applicable risks and uncertainties include the risks and uncertainties, among other things, regarding: the success in the development and potential commercialization of our product candidates, including defactinib and other compounds in combination with avutometinib (VS-6766); the occurrence of adverse safety events and/or unexpected concerns that may arise from additional data or analysis or result in unmanageable safety profiles as compared to their levels of efficacy; or our ability to obtain, maintain and enforce patent and other intellectual property protection for our product candidates.

Other risks and uncertainties include those identified under the heading "Risk Factors" in the Company's Annual Report on Form 10-K for the year ended December 31, 2022, as filed with the Securities and Exchange Commission (SEC) on March 14, 2023, and in any subsequent filings with the SEC, which are available at www.sec.gov and www.verastem.com.

The forward-looking statements in this presentation speak only as of the original date of this presentation, and we undertake no obligation to update or revise any of these statements.

Verastem Oncology

Well Positioned to Capitalize on Growth Opportunities

Lead clinical program has best-in-class potential

Avutometinib (VS-6766; RAF/MEK clamp) and defactinib (FAK inhibitor) are clinically active against RAS pathway-driven cancers

Rapid development path to market in LGSOC

FDA Breakthrough Therapy Designation; Interim analysis of Part A of registration-directed RAMP 201 trial: avutometinib + defactinib confirmed response rates and safety profile support continued development in all recurrent LGSOC; timing of accelerated approval filing to be based on data maturity and finalization of confirmatory study plans

Significant downstream market opportunity and blockbuster potential

30% of all human cancers are driven by mutations in RAS; Avutometinib combinations potentially broadly applicable across a variety of tumor types.

Clinical collaborations with Amgen & Mirati evaluating the combinations of avutometinib with sotorasib & adagrasib, respectively, in KRAS G12C NSCLC supported by strong pre-clinical rationale

Multiple clinical studies in progress evaluating avutometinib combinations across RAS pathway-driven cancers

Patent Update

Recently issued intermittent dosing IP for both avutometinib alone and avutometinib + defactinib extends patent coverage up to 2038 and 2040

Strong balance sheet

Cash balance of \$87.9 million as of December 31, 2022

Pro-forma cash balance of \$132.9 million including Q1 2023 financing activities (\$30 million from Series B Preferred Stock issuance and \$15 million from Oxford Term B Loan draw)

Up to \$150 million of non-dilutive funding available from credit facility

Company ended Q4 2022 with \$15.6 million non-GAAP operating expenses*

We are a biopharmaceutical company committed to developing and commercializing new medicines for patients battling cancer

* Q4 2022 GAAP operating expenses - \$16.85M minus Q4 2022 stock compensation of \$1.29M = \$15.56M Q4 2022 non-GAAP operating expenses

Key VSTM Achievements 2022 & Anticipated Milestones 2023





	1H2022	2H2022	1Q2023	2Q2023	2H2023
LGSOC	✓ RAMP 201 Selection Phase Complete; Initiated enrollment of Expansion Phase ✓ RAMP 201 Selection Phase Update** ✓ Translational data from FRAME LGSOC cohort presented	✓ RAMP 201 Second Interim Update ✓ RAMP 201 FDA Meeting - Avuto + defactinib selected as Go-Forward	✓ RAMP 201 Complete target enrollment of Expansion Phase ✓ Launch LGSOC patient education campaign	- Present updated results of Part A RAMP 201 - Discuss confirmatory trial study design for recurrent LGSOC program	Initiate confirmatory study of avutometinib + defactinib in recurrent LGSOC
NSCLC	✓ RAMP 202 Complete enrollment of Selection Phase ✓ Initiate RAMP 203 (avuto + sotorasib) G12C ✓ Top-Line Data from avutometinib + everolimus in KRAS mt ✓ RAMP 202 Amended to include BRAF mt cohorts	✓ Initiate RAMP 204 (avuto + adagrasib) G12C ✓ Top-Line Data from RAMP 202 Selection Phase ✓ RAMP 203 advanced to final dose level		- RAMP 203: Determine recommended phase 2 dose - Present updated results of IST avutometinib + everolimus in KRAS mt NSCLC	- RAMP 203: Report initial read-out of safety and preliminary efficacy - RAMP 204: Initial read-out of safety and recommended dose
Additional Indications	✓ PanCAN Agreement Executed	✓ Initiate combo study of avutometinib + cetuximab in KRAS mt CRC * ✓ Initiate RAMP 205 combo avutometinib + gemcitabine/nab-paclitaxel + defactinib in metastatic pancreatic cancer	- Initiate combo study of avutometinib + pembrolizumab in BRAF mt melanoma * - Initiate thyroid cancer *	- Pediatric Cancer * - Early safety data of avutometinib + cetuximab in KRAS mt CRC * - AACR: Everolimus NSCLC presentation Preclinical-combinations presentations *	- RAMP 205: determine recommended phase 2 dose and complete enrollment of initial phase 2 expansion cohort - Initial results of Gynecological basket trial * - Early efficacy results of thyroid cancer * - Early safety results of avutometinib + pembrolizumab in BRAF melanoma *

Avutometinib is a Differentiated Agent with the Potential to Serve as the Backbone for Combinations Across RAS Pathway-Driven Cancers

- Unique RAF/MEK clamp mechanism of action
- Novel intermittent dosing schedule; convenient oral regimen
- Breakthrough Therapy Designation in recurrent low-grade serous ovarian cancer(LGSOC) regardless of KRAS status
- Potential best-in-class safety & tolerability profile relative to marketed MEK inhibitors, with potential for combinability with agents from multiple target classes
- Promising signals of clinical activity in various RAS pathway-driven cancers, including in patients whose tumors previously progressed on other MEK inhibitors
- Preclinical anti-proliferative activity across multiple MAPK pathway alterations (e.g. KRAS, NRAS, BRAF, NFI mt) and multiple solid tumor indications
- Strong preclinical combination data with other agents targeting RAS pathway (e.g. KRAS G12C inhibitors) and parallel pathways (e.g. FAK inhibitors)

Strong Scientific Rationale for Avutometinib and FAK Inhibitor Combination

Anti-Tumor Efficacy in KRAS Mutant and Wild-Type LGSOC models




LGSOC Unmet Need & Opportunity

- LGSOC is a less common type of ovarian cancer that is often diagnosed in younger women
 - LGSOC is a unique disease that is distinct from high-grade serous ovarian cancer (HGSOC) in its pathology, protracted clinical course and low response to chemotherapy and thus requires a more tailored therapeutic approach
 - An estimated 1,000-2,000 patients are diagnosed with LGSOC per year in the U.S., with prevalence of ~6,000
- There are currently no approved therapies specifically indicated for recurrent LGSOC
 - Recent clinical trials in recurrent LGSOC showed that standard-of-care chemo and endocrine therapy are relatively ineffective (6-13% ORR).
 - Currently available therapies have a poor benefit/risk profile and NCCN guidelines include clinical trials and observation highlighting the lack of approved & effective therapies
- LGSOC is known to be driven by the MAPK (RAS) pathway in $\geq 70\%$ of patients
- A phase I/II study in the UK (FRAME) evaluated the combination of avutometinib and defactinib
 - Results in recurrent LGSOC showed a 46% confirmed ORR with durable responses and favorable safety/tolerability

- Breakthrough Therapy Designation was granted for avutometinib and defactinib in recurrent LGSOC
- RAMP 201: A registration-directed Phase 2 trial of avutometinib + defactinib in recurrent LGSOC is ongoing

RAMP 201 (ENGOTov60/GOG3052): Registration-Directed Phase 2 Trial of Avutometinib ± Defactinib in Patients with Recurrent LGSOC




*Dosing: Avutometinib + Defactinib combo: Avutometinib 3.2 mg PO 2x/wk 21/28 days + Defactinib 200 mg PO BID: 21/28 days;
 Avutometinib monotherapy: Avutometinib 4.0 mg PO 2x/wk 21/28 days

**Expansion Phase – Final sample size to be adjusted based on adaptive design

Combination Arm:
 ❖ Expanded Enrollment Ongoing
 ✓ Target Enrollment Reached (N=72)

RAMP 201 (ENGOTov60/GOG3052): Registration-Directed Phase 2 Trial of Avutometinib ± Defactinib in Patients with Recurrent LGSOC



Planned Interim Analysis:

- ❑ **Initial Efficacy Data: Part A (Evaluable Patients)**
- ❑ **Safety Data: All Treated Patients**

- **Heavily** pretreated population;
 - Median of **4** prior lines of systemic therapy
- Clinical development continues in **all recurrent LGSOC**, regardless of KRAS status
- **Combination of avutometinib and defactinib** selected as “Go Forward” treatment regimen
 - **ORR 28%** independently confirmed
 - KRAS mt: 27%; KRAS wt: 29%
 - Tolerable safety profile - No new safety signals;
 - Majority of evaluable patients remain on treatment (62%)

Efficacy and Tolerability of Avutometinib and Defactinib in LGSOC

A Head-to-Head Study Has Not Been Performed

Trial	Median Number of Prior lines of Therapy	Prior MEK Allowed	Prior Bevacizumab	Therapy	Response Rate ORR	Image Assessment	Median PFS Months (95% CI)	Discontinuation Rate Due to AEs
GOG 281 ¹	3 (1-10)	No	* Low %	Standard of Care	6% ^	INV	7.2 (5.6-9.9)	13%
				Trametinib	26%^	INV	13.0 (9.9-15.0)	36%
MILO ²	2 (1-8)	No	* Low %	Standard of Care	13%	BICR	10.6 (9.2 to 14.5)	17%
				Binimetinib	16%	BICR	9.1 (7.3-11.3)	31%
FRAME ³	3	Yes	12 %	Avutometinib + Defactinib	46%^ 95% CI: (26%, 67%)	INV	23 (11 - NR)	4%
RAMP 201 Part A ³	4	Yes	65%	Avutometinib + Defactinib	28% 95% CI: (13%, 47%) 38%**	BICR	Not Yet Available	9%

* Low historical use of bevacizumab during trial conduct.

** Confirmed + Unconfirmed Objectives responses

^ Investigator Assessment only (No independent Central Review of Imaging)

Standard of Care = letrozole, tamoxifen, chemotherapy

INV = Investigator

BICR = Blinded independent central review

PFS = Progression free survival

³ Avutometinib + Defactinib combination evaluable population

RAMP 201: Safety and Tolerability Profile of Avutometinib + Defactinib

No New Safety Signals; Few Discontinuations Due to Adverse Events

Most Common Treatment-Related Adverse Events (>20%) in All Treated Patients

- Majority of adverse events are mild to moderate
- Adverse events are monitorable, manageable and reversible
- Few discontinuations due to adverse events (9% in combo n=5 – includes 3 pts for elevated blood CPK laboratory values*)


* No association to date with clinically significant toxicities, including rhabdomyolysis

Avutometinib + Defactinib						
	Any Grade			Grade ≥3		
	KRAS mt (n=22) %	KRAS wt (n=35) %	Total (n=57) %	KRAS mt (n=22) %	KRAS wt (n=35) %	Total (n=57) %
Nausea	45.5	65.7	57.9	0	0	0
Diarrhea	54.5	51.4	52.6	4.5	2.9	3.5
Edema peripheral	36.4	48.6	43.9	0	0	0
Fatigue	40.9	40.0	40.4	9.1	2.9	5.3
Vision blurred	31.8	42.9	38.6	0	0	0
Vomiting	40.9	37.1	38.6	0	0	0
Blood creatine phosphokinase increased	40.9	37.1	38.6	18.2	14.3	15.8
Dermatitis acneiform	27.3	45.7	38.6	4.5	0	1.8
Rash	31.8	37.1	35.1	0	2.9	1.8
Dry skin	13.6	31.4	24.6	0	0	0

RAMP 201 Part A Results and FDA Feedback Establish Path to Filing for Accelerated Approval

Update

- Combination of avutemetinib with defactinib selected as go forward treatment regimen
- Combination development continues in all recurrent LGSOC, regardless of KRAS status
- Encouraging efficacy results include independently confirmed responses
- No new safety signals; few discontinuations due to adverse events
- Majority of patients remain on treatment



Next Steps

- Target enrollment for primary analysis (n=72) in combination has been achieved
- Plan to file for accelerated approval based on the totality of the data from the RAMP 201 and FRAME studies
- Continued enrollment in RAMP 201 combination arm only is planned to expand clinical experience in anticipation of initiation of a confirmatory study
- The Company will provide an update after agreement with the FDA on the confirmatory study
- The Company is planning a RAMP 201 presentation at a scientific conference in mid-2023

Comprehensive approach to establish more complete blockade of RAS pathway & resistance pathways

	Indication	Incidence/ Prevalence	Regimen	Setting
Gynecologic Cancers	LGSOC	Prevalence ¹ : 6K 	Avutometinib + defactinib	Relapsed Refractory molecularly profiled LGSOC
	Gynecologic Basket*	Incidence ⁶⁻¹⁰ : 85K	Avutometinib + defactinib	Recurrent RAS Pathway-driven (RAS/RAF/NF1) endometrioid cancer, mucinous ovarian cancer, high-grade serous ovarian cancer or cervical cancer
NSCLC Adenocarcinoma	KRAS G12C	Incidence ^{2,3} : 114K 	Avutometinib + sotorasib  Avutometinib + adagrasib 	Recurrent KRAS G12C with prior KRAS G12C inhibitor(i) treatment or KRAS G12Ci naïve Recurrent KRAS G12C with prior KRAS G12Ci treatment that progressed
	BRAF mt	Incidence ^{2,3} : 114K 	Avutometinib + defactinib	Recurrent BRAFV600E & non-V600E mutant NSCLC
Pancreatic	PDAC	Incidence ⁴ : 58K 	Avutometinib + defactinib + gemcitabine/nab-paclitaxel 	Previously untreated (front-line) metastatic pancreatic ductal adenocarcinoma (PDAC)
CRC	KRAS mt*	Incidence ⁵ : 148K 	Avutometinib + cetuximab	Recurrent metastatic KRAS mt
Breast Cancer	ER+*	Incidence ⁵ : 279K 	Avutometinib + abemaciclib + fulvestrant	Recurrent ER+/HER2- breast cancer following progression on CDK4/6i + aromatase inhibitor
Melanoma	BRAFV600E*	Incidence ⁴ : 108K 	Avutometinib + pembrolizumab	Recurrent BRAFV600E/K or NRAS (Phase I only) mutant Melanoma following progression on prior anti-PD1 therapy

*IST

¹ References: Monk, Randall, Grisham, The Evolving Landscape of Chemotherapy in Newly Diagnosed Advanced Epithelial Ovarian Cancer, Am Soc Clin Oncol Educ Book; 2019; Slomovitz, Gourley, Carey, Malpica, Shih, Huntsman, Fader., Grisham et al, Low-Grade serous ovarian cancer: State of the Science; Gynecol Oncol; 2020. Grisham, Iyer, Low-Grade Serous Ovarian Cancer: Current Treatment Paradigms and Future Directions; Curr Treat Options Oncology; 2018; Globocan 2020; ²Pakkala and Ramalingam JCI Insight 2018); ³Cancer Statistics 2020, Siegel et. al. CA Cancer J Clin 2020;70:7-30; ⁴Cancer Statistics 2020, Siegel et. al. CA Cancer J Clin 2020;70:7-30 ⁵CbioPortal; ⁶Uterine cancer is one of the leading gynecologic neoplastic disorders in the US, of which over 80% are endometrioid adenocarcinomas (EA); ⁷Endometrioid OC (EnOC) accounts for approximately 10% of all OC, with the majority of cases diagnosed as low grade, early stage disease with excellent clinical; ⁸Mucinous ovarian cancer: 3-11% of ovarian cancer (Hada et al., 2021); ⁹90% of Ovarian Cancer is Epithelial Ovarian Cancer (<https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2018/cancer-facts-and-figures-2018.pdf>); ¹⁰HGSOC the most common type of ovarian cancer, accounting for approximately 75% of epithelial ovarian cancers. (<https://ocrahope.org/news/high-grade-serous-carcinoma/#:~:text=High%2Dgrade%20serous%20carcinoma%20is,unless%20another%20type%20is%20specified.>)

Broad Development Opportunities Across Multiple RAS/MAPK Pathway-Driven Cancers


High Priority Registration Indication

Registration-Directed Trial Initiated in 4Q20

- LGSOC^{1,2} (RAMP 201)-Target enrollment reached

RAS Pathway Dependent Cancers

- Gynecological^{1,2}
- NSCLC^{1,2}
- Colorectal^{1,2}
- Melanoma^{1,2}
- Pancreatic²
- Thyroid^{1,2}



Key Signal Finding

- Avutometinib + G12Ci in KRAS G12C NSCLC² (RAMP 203 - sotorasib) & (RAMP 204 - adagrasib)
- Avutometinib + defactinib in BRAF mt (V600E & non-V600E) NSCLC^{1,2} (RAMP 202)
- Avutometinib + defactinib and gemcitabine/nab-paclitaxel in first line pancreatic cancer (RAMP 205)²
- Avutometinib + defactinib in RAS/RAF/NFI mt gynecological cancers^{1,2}
- Avutometinib + cetuximab in KRAS mt CRC²
- Avutometinib + abemaciclib and fulvestrant in ER+ breast cancer²
- Avutometinib + pembrolizumab in BRAFV600E melanoma²

Rational Combinations

- KRAS inhibitors² (G12Ci & G12Di)
- Anti-EGFR²
- Everolimus^{1,2}
- CDK4/6 inhibitor²
- Anti-PD-1^{1,2}
- Chemotherapy²

Biomarker Selection

- KRAS mt^{1,2}
- BRAF mt (V600 & non-V600)^{1,2}
- NRAS mt^{1,2}
- CRAF mt/fusions²

Robust Clinical Program: Avutometinib in multiple combinations across RAS/MAPK pathway-driven tumors

INDICATION	REGIMEN	STUDY NAME	PRECLINICAL	PHASE I	PHASE 2	PHASE 3	CLINICAL COLLABORATION WITH
LGSOC ¹	Avutometinib + defactinib	RAMP 201	Registration-directed trial: accelerated approval cohort fully enrolled				
R/R LGSOC	Avutometinib + defactinib	IST-FRAME					
Gynecological Cancers (RAS Pathway-driven) ²	Avutometinib + defactinib	IST					
Mesonephric ²	Avutometinib + defactinib	IST					
R/R NSCLC (BRAF mt)	Avutometinib + defactinib	RAMP 202					  
R/R NSCLC (KRAS G12C)	Avutometinib + sotorasib	RAMP 203					
R/R NSCLC (KRAS G12C)	Avutometinib + adagrasib	RAMP 204					
Pancreatic Ductal Adenocarcinoma	Avutometinib + gemcitabine/nab-paclitaxel + defactinib	RAMP 205					
R/R NSCLC (KRAS mt)	Avutometinib + everolimus (mTORi)	IST					
R/R Colorectal Cancer (KRAS mt)	Avutometinib + cetuximab (EGFRi)	IST					
ER+ Breast Cancer	Avutometinib + abemaciclib + fulvestrant	IST					
BRAF V600E Melanoma ²	Avutometinib + pembrolizumab	IST					

Key Financial Statistics

As of and for the quarter ended December 31, 2022

Cash, cash equivalents & investments	\$87.9M
Pro-forma cash, cash equivalents, & investments	\$132.9M [^]
Non-GAAP Operating Expenses*	\$15.6M
Shares Outstanding	200.5M

[^] Pro-forma cash, cash equivalents & investments includes Q1 2023 financing activities (\$30M from Series B Preferred Stock issuance and \$15M from Oxford Term B Loan draw).


Sources of Non-Dilutive Capital

- **Oxford Finance LLC Credit Facility**
 - Up to \$150M available in a series of term loans
 - \$40M term loans outstanding as of March 2023.
 - Remaining \$110M available upon achievement of pre-defined milestones or at lender's discretion
 - Floating interest rate, subject to a floor and a cap; 5% final payment charge, and loan subject to 1-3% early payment fee
 - Interest only payments through April 2025
 - No financial covenants
- **Secura Bio, Inc. (Secura) Asset Purchase Agreement – COPIKTRA**
 - Regulatory and commercial milestone payments up to \$95M
 - Entitled to receive 50% of royalties, milestones, and sublicensee revenue payments made to Secura related to COPIKTRA
 - Low double-digit royalties on annual net sales over \$100M in US, EU, and UK




Avutometinib RAF/MEK Clamp Program Overview

The background of the slide is a solid blue color. On the right side, there are several diagonal stripes in white, teal, and orange. A horizontal teal stripe runs across the bottom of the slide, partially overlapping the diagonal stripes.

Avutometinib is a Unique Small Molecule RAF/MEK Clamp






- Avutometinib inhibits MEK, BRAF & CRAF by trapping these molecules in inactive complexes
- MEK inhibitors paradoxically induce MEK phosphorylation (pMEK) by relieving ERK-dependent feedback inhibition of RAF
- By inhibiting RAF phosphorylation of MEK, avutometinib has advantage of not inducing pMEK
- Avutometinib inhibits ERK signaling more completely; may confer enhanced therapeutic activity




Avutometinib is a Unique RAF/MEK Clamp which Induces Inactive Complexes of MEK with ARAF, BRAF & CRAF

Contrasting mechanism of action vs. trametinib



Avutometinib Inhibits Cell Proliferation Across Multiple RAS/MAPK Pathway Alterations and Multiple Solid Tumor Histologies



Optimized Dosing Schedule Defined: Favorable Tolerability Profile with Novel Intermittent Dosing Regimen

Summary of Adverse Events Grade ≥ 3 Occurring in $\geq 5\%$ of patients

	Avutometinib monotherapy Daily at MTD N=6 28-day cycle	RP2D Avutometinib monotherapy 4mg twice weekly N=26 28-day cycle	RP2D (Avutometinib 3.2mg twice weekly + defactinib 200mg twice daily) N=38 21 days of 28-day cycle
Treatment Related Adverse Event	Grade ≥ 3	Grade ≥ 3	Grade ≥ 3
Rash	3 (50%)	5 (19%)	2 (5%)
CK elevation (Creatine phosphokinase)	1 (17%)	2 (8%)	2 (5%)

Avutometinib Patent Exclusivity

Composition of Matter

PTE

Feb 2027 + 5 yrs (PTE) = 2032

Method of Making

Sept 2032

Dosing Protocol

May 2038

Combination w/ Defactinib

Sept 2040


Methods of Treating

2041 - 2042 if issued

Combinations

2041 - 2042 if issued

Avutometinib ± Defactinib in Low-Grade Serous Ovarian Cancer

The background of the slide features a large blue area on the left. On the right side, there are several diagonal stripes in white, teal, and orange. At the bottom, there is a horizontal teal stripe and a blue stripe.

LGSOC is a Unique RAS Pathway-Driven Cancer with a High Unmet Need



LGSOC is a type of ovarian cancer that disproportionately affects younger women



1,000 to 2,000 patients in the U.S. and 15,000 to 30,000 worldwide diagnosed with LGSOC each year



A slow growing cancer, that has a median survival of almost 10 years, so patients remain in treatment for a long time (10-yr prevalence ~80,000 worldwide, ~6,000 US)




Patients often experience significant pain and suffering from their disease over time



Most prior research has focused on high grade serous ovarian cancer (HGSOC). However, LGSOC is clinically, histologically and molecularly unique from HGSOC with limited treatments available


~30% of LGSOC Patients Have KRAS mt
~70% of LGSOC Shows RAS Pathway-Associated mts




References: AACR Project GENIE Cohort v9.0-public and Verastem unpublished analysis

Low-Grade and High-Grade Serous Ovarian Cancer Are Different Diseases

Variable	LGSOC	HGSOC
Nuclear atypia	Uniform round to oval with little variation	+++ Marked variation
Mitotic Index	<12 mitoses per 10 hpf	>12 mitoses per 10 hpf
Chromatin and variation in size of nucleus	Little	Marked (nuclear size ratio ≥ 3)
Mutation	KRAS ++ BRAF + ER/PR +++	P53 +++ BRCA1/2 +
Precursor	Serous borderline tumor	Tubal intraepithelial neoplasia



LGSOC




HGSOC

Recurrent LGSOC: High Medical Need

No Approved Treatment Options – Limited Benefit from Available Therapies

Recurrent Low-Grade Ovarian Cancer – Treatment Guidelines ¹



Recent LGSOC Trials Highlight High Unmet Need

Trial	Number of Prior Systemic Therapies Median (range)	Prior MEK allowed	Prior Bevacizumab	Therapy	Response Rate ORR	Image Assessment	Median PFS Months (95% CI)	Discontinuation Rate due to AEs
GOG 281 ¹	3 (1-10)	No	* Low %	SoC	6%	INV	7.2 (5.6-9.9)	13%
				Trametinib	26%	INV	13.0 (9.9-15.0)	36%
MILO ²	2 (1-8)	No	* Low %	SoC	13%	BICR	10.6 (9.2 - 14.5)	17%
				Binimetinib ²	16%	BICR	9.1 (7.3-11.3)	31%

¹ Study GOG 281 trial Gershenson et al., Lancet 2022

² MILO Study Monk et al., J Clin Oncol 2020.

* Low historical use of bevacizumab during trial conduct. % not reported

MILO: no more than 3 lines of prior chemotherapy

SoC = Standard of Care
(endocrine / chemotherapy)

INV = Investigator

BICR = Blinded independent central review


PFS = Progression free survival

CI = confidence interval

NR = Not reached


FRAME Study: Solid Foundation for the Development of Avutometinib in Combination with Defactinib in Recurrent LGSOC (n=24)

Response by RECIST



- Overall response rate (ORR) = 46% (11 confirmed PRs/24)
 - KRAS mutant ORR = 64% (7 confirmed PRs/11)
 - KRAS wild-type ORR = 44% (4 confirmed PRs/9)
 - KRAS status undetermined (1 unconfirmed PR/4)
- Response too early to determine for 2 pts on study for ≤ 5 months
- Median 3 lines of Prior Treatment (Prior MEKi 10 pts, Prior Bev 4 pts)
- Responses in patients previously treated with MEKi
- 54% (13/24) patients still on treatment
- 1 patient discontinuing for adverse events as of April 2021
- Median PFS 23 months (95% CI 10.6-NR) across all LGSOC

Best response by RECIST



Data cut off April 2021


MEKi: MEK inhibitor

Bev: Bevacizumab

PFS: Progression free survival

NR: Not reached

RAMP 201 (ENGOTov60/GOG3052): Registration-Directed Phase 2 Trial of Avutometinib ± Defactinib in Patients with Recurrent LGSOC




*Dosing: Avutometinib + Defactinib combo: Avutometinib 3.2 mg PO 2x/wk 21/28 days + Defactinib 200 mg PO BID: 21/28 days;
 Avutometinib monotherapy: Avutometinib 4.0 mg PO 2x/wk 21/28 days

**Expansion Phase – Final sample size to be adjusted based on adaptive design

Combination Arm:
 ❖ Expanded Enrollment Ongoing
 ✓ Target Enrollment Reached (N=72)

RAMP 201 (ENGOTov60/GOG3052): Registration-Directed Phase 2 Trial of Avutometinib ± Defactinib in Patients with Recurrent LGSOC



Planned Interim Analysis:

- ❑ **Initial Efficacy Data: Part A (Evaluable Patients)**
- ❑ **Safety Data: All Treated Patients**

- **Heavily** pretreated population;
 - Median of **4** prior lines of systemic therapy
- Clinical development continues in **all recurrent LGSOC**, regardless of KRAS status
- **Combination of avutometinib and defactinib** selected as “Go Forward” treatment regimen
 - **ORR 28%** by blinded independent central review
 - KRAS mt: 27%; KRAS wt: 29%
 - Tolerable safety profile - No new safety signals;
 - Majority of evaluable patients remain on treatment (62%)

RAMP 201 Part A: Heavily Pre-Treated Patient Population

*Prior Platinum-Based Chemotherapy, Endocrine Therapy, Bevacizumab in Most Patients;
Prior MEK Inhibitor Therapy was Permitted*

	Avutometinib Monotherapy			Avutometinib + Defactinib		
	KRAS mt (n=16)	KRAS wt (n=17)	Total (n=33)	KRAS mt (n=16)	KRAS wt (n=15)	Total (n=31)
Age (yrs), median (min, max)	58 (27, 72)	48 (27, 74)	51 (27, 74)	61 (29, 71)	50 (30, 74)	55 (29, 74)
ECOG, n (%)						
0	8 (50)	15 (88)	23 (70)	11 (69)	9 (60)	20 (65)
1	8 (50)	2 (12)	10 (30)	5 (31)	6 (40)	11 (35)
Number of Prior Systemic Regimens, median (min, max)	3.0 (1, 10)	3.0 (1, 9)	3.0 (1, 10)	4.0 (1, 9)	4.0 (1, 11)	4.0 (1, 11)
Prior platinum-based chemotherapy, n (%)	15 (93.8)	17 (100)	32 (97.0)	16 (100)	15 (100)	31 (100)
Prior MEK inhibitor therapy, n (%)	5 (31.3)	5 (29.4)	10 (30.3)	2 (12.5)	2 (13.3)	4 (12.9)
Prior Bevacizumab, n (%)	8 (50.0)	8 (47.1)	16 (48.5)	7 (43.8)	13 (86.7)	20 (64.5)
Prior Hormonal therapy, n (%)	11 (68.8)	12 (70.6)	23 (69.7)	15 (93.8)	13 (86.7)	28 (90.3)

RAMP 201 Part A: Evaluable Patient Population*

Positive ORR Confirmed by Blinded Independent Central Review (BICR) Support
Avutometinib + Defactinib as Go Forward Regimen in LGSOC - Regardless of KRAS Status

	Avutometinib			Avutometinib + Defactinib		
	KRAS mt (n=15)	KRAS wt (n=15)	Total (n=30)	KRAS mt (n=15)	KRAS wt (n=14)	Total (n=29)
ORR, n(%)						
Confirmed PR	2 (13)	0	2 (7)	4 (27)	4 (29)	8 (28) 95% CI: (13%, 47%)
Confirmed + Unconfirmed PR**	2 (13)	0	2 (7)	7 (47)	4 (29)	11 (38)
SD**	12 (80)	13 (87)	25 (83)	11 (73)	8 (57)	19 (66)
Disease Control Rate	14 (93)	13 (87)	27 (90)	15 (100)	12 (86)	27 (93%)
PD	1 (7)	2 (13)	3 (10)	0	2 (14)	2 (7)

* Evaluable for Efficacy: At least one blinded imaging assessment in 30 of 33 and 29 of 31 patients enrolled in respective treatment arms

** Includes patients with unconfirmed PR who have a chance to be confirmed at their next assessment

Go Forward Regimen: Combination of Avutometinib and Defactinib

High Disease Control Rate + Tumor Reduction in Recurrent LGSOC

Part A (Evaluable for Efficacy *)


Confirmed ORR: **28%**

Confirmed/Unconfirmed ORR: **38%**

Disease Control Rate (SD+PR): **93%**

Patients still on study treatment: **62%**


Shortest follow up: 5 months




Go Forward Regimen: Combination of Avutometinib and Defactinib

Initial Data from RAMP 201 Trial Reinforce Findings from FRAME Trial

RAMP 201 (Part A)
Interim Analysis - Blinded ICR
N=29



FRAME
Investigator Assessment
N=24



RAMP 201: Safety and Tolerability Profile of Avutometinib + Defactinib

No New Safety Signals; Few Discontinuations Due to Adverse Events

Most Common Treatment-Related Adverse Events (>20%) in All Treated Patients

- Majority of adverse events are mild to moderate
- Adverse events are monitorable, manageable and reversible
- Few discontinuations due to adverse events (9% in combo n=5 – includes 3 pts for elevated blood CPK laboratory values*)


* No association to date with clinically significant toxicities, including rhabdomyolysis

Avutometinib + Defactinib						
	Any Grade			Grade ≥3		
	KRAS mt (n=22) %	KRAS wt (n=35) %	Total (n=57) %	KRAS mt (n=22) %	KRAS wt (n=35) %	Total (n=57) %
Nausea	45.5	65.7	57.9	0	0	0
Diarrhea	54.5	51.4	52.6	4.5	2.9	3.5
Edema peripheral	36.4	48.6	43.9	0	0	0
Fatigue	40.9	40.0	40.4	9.1	2.9	5.3
Vision blurred	31.8	42.9	38.6	0	0	0
Vomiting	40.9	37.1	38.6	0	0	0
Blood creatine phosphokinase increased	40.9	37.1	38.6	18.2	14.3	15.8
Dermatitis acneiform	27.3	45.7	38.6	4.5	0	1.8
Rash	31.8	37.1	35.1	0	2.9	1.8
Dry skin	13.6	31.4	24.6	0	0	0

RAMP 201 Part A Results and FDA Feedback Establish Path to Filing for Accelerated Approval

Update

- Combination of avutometinib with defactinib selected as go forward treatment regimen
- Combination development continues in all recurrent LGSOC, regardless of KRAS status
- Encouraging efficacy results include independently confirmed responses
- No new safety signals; few discontinuations due to adverse events
- Majority of patients remain on treatment



Next Steps

- Target enrollment for primary analysis (n=72) in combination has been achieved
- Plan to file for accelerated approval based on the totality of the data from the RAMP 201 and FRAME studies
- Continued enrollment in RAMP 201 combination arm only is planned to expand clinical experience in anticipation of initiation of a confirmatory study
- The Company will provide an update after agreement with the FDA on the confirmatory study
- The Company is planning a RAMP 201 presentation at a scientific conference in mid-2023

Efficacy and Tolerability of Avutometinib and Defactinib in LGSOC

A Head-to-Head Study Has Not Been Performed

Trial	Median Number of Prior lines of Therapy	Prior MEK Allowed	Prior Bevacizumab	Therapy	Response Rate ORR	Image Assessment	Median PFS Months (95% CI)	Discontinuation Rate Due to AEs
GOG 281 ¹	3 (1-10)	No	* Low %	Standard of Care	6% ^	INV	7.2 (5.6-9.9)	13%
				Trametinib	26%^	INV	13.0 (9.9-15.0)	36%
MILO ²	2 (1-8)	No	* Low %	Standard of Care	13%	BICR	10.6 (9.2 to 14.5)	17%
				Binimetinib	16%	BICR	9.1 (7.3-11.3)	31%
FRAME ³	3	Yes	12 %	Avutometinib + defactinib	46%^ 95% CI: (26%, 67%)	INV	23 (11 - NR)	4%
RAMP 201 Part A ³	4	Yes	65%	Avutometinib + defactinib	28% 95% CI: (13%, 47%) 38%**	BICR	Not Yet Available	9%

* Low historical use of bevacizumab during trial conduct.

** Confirmed + Unconfirmed Objectives responses

^ Investigator Assessment only (No independent Central Review of Imaging)

Standard of Care = letrozole, tamoxifen, chemotherapy

INV = Investigator

BICR = Blinded independent central review

PFS = Progression free survival

CI = confidence interval



NR = Not reached

³ Avutometinib + Defactinib combination evaluable population

¹ Study GOG 281 trial Gershenson et al., Lancet 2022

² MILO Study Monk et al., J Clin Oncol 2020.

RAMP 201 Part A Interim Data Support Meaningful Market Potential for All Recurrent LGSOC Regardless of KRAS Status with Long Duration of Therapy



¹ References: Monk, Randall, Grisham, The Evolving Landscape of Chemotherapy in Newly Diagnosed Advanced Epithelial Ovarian Cancer, Am Soc Clin Oncol Educ Book; 2019; Slomovitz, Gourley, Carey, Malpica, Shih, Huntsman, Fader., Grisham et al, Low-Grade serous ovarian cancer: State of the Science; Gynecol Oncol; 2020. Grisham, Iyer, Low-Grade Serous Ovarian Cancer: Current Treatment Paradigms and Future Directions; Curr Treat Options Oncology; 2018; Globocan 2020

² Patient-months of Therapy metric calculated by multiplying relevant incidence/prevalence rate times estimated duration of therapy; represents US market opportunity only; patient population estimates from Globocan 2020, American Cancer Society 2021, AACR Genie Cohort V9.0 public, and scientific publications. Duration of therapy estimates from clinical studies and clinician experience. Patient-months on therapy is for 2nd-line+ patients

³ NSCLC KRAS G12C 2nd line patients (incidence); Pancreatic RAS/RAF mutant 2nd-line patients (incidence); LGSOC KRAS mutant and wild-type patients (prevalence); Endometrioid RAS/RAF mutant 2nd-line patients (incidence); Uveal melanoma RAS/RAF mutant 2nd-line patients (incidence)

RAS Pathway-Driven Cancers and Rational Avutometinib Combinations

The background of the slide features a series of parallel diagonal stripes in blue, teal, and orange, creating a modern, geometric design.

High Unmet Needs in Additional RAS/MAPK Pathway-Driven Cancers




KRAS-mutant Cancers¹


NSCLC
Incidence^{3,5}:
114K




Pancreatic
Incidence⁵:
58K



Uterine Endometrioid
Incidence^{4,5}: 59K



Colorectal
Incidence⁵:
148K




NRAS-mutant Cancers¹

Melanoma
Incidence⁵:
108K



Multiple Myeloma
Incidence⁵:
32K



BRAF-mutant Cancers²


Melanoma
Incidence⁵:
108K



Colorectal
Incidence⁷:
148K



Papillary Thyroid
Incidence^{5,6}:
42K



NSCLC
Incidence^{3,7}:
194K



Breadth of potential opportunity

- 30% of all human cancers are driven by mutations of the RAS family of genes⁶

Established prognostic significance

- Patients with mutations of the RAS family have an overall worse prognosis

Challenges with conventional approaches

- Modest progress; limited number of approved therapies
- Single agent therapies (e.g., MEK inhibitors) associated with resistance
- Tolerable combination regimens with MEK inhibitors have been challenging
- Approved RAS inhibitors address only a minority of all RAS mutated cancers (KRAS G12C)

Incidence References:


¹Reference for RAS mt frequencies – Cox et al. *Nature Reviews* 13: 828, 2014; ²Reference for BRAF mt frequencies – Turski et al. *Mol Cancer Ther* 15: 533, 2016

³50% of NSCLC is adenocarcinoma (Pakkala and Ramalingam *JCI Insight* 2018); ⁴90% of all uterine cancers are of the endometrial type (ACS); ⁵Cancer Statistics 2020, Siegel et. al. *CA Cancer J Clin* 2020;70:7-30; ⁶8 out of 10 thyroid cancers are of the papillary type (ACS)⁷CbioPortal

References:

McCormick F *Clin Cancer Res* 15April2015; ⁶Adderley H et al. *EBioMedicine* 01Mar2019; Papke B et al. *Science* 17Mar2017; Ryan M et al. *Nature Reviews Clinical Oncology* 01Oct2018; NIH cancer.gov/research/key-initiatives/ras

Vertical Blockade: Establishing Avutometinib as the Backbone of Therapy for RAS/MAPK Pathway-Driven Tumors




■ Current Challenges

- Blocking any single target in the pathway is insufficient for maximum depth and duration of anti-tumor efficacy
 - e.g. SHP2i, KRAS-G12Ci, KRAS-G12Di, RAFi, MEKi, ERKi
- Vertical blockade concept is now well established
 - Necessary to block more than 1 target in the pathway
- Many of these agents (e.g. SHP2i, MEKi) have poor tolerability as monotherapy and in combination

■ Solutions offered by Avutometinib

- Vertical blockade (RAF and MEK blockade) in a single drug
- Potential best-in-class tolerability with recommended twice weekly dosing regimen
 - Should enable tolerable combinations
- Compelling synergy data (preclinical) for avutometinib combinations (e.g. with KRAS G12C inhibitors) supporting clinical combinations
- Ongoing clinical combination studies with G12Ci (sotorasib, adagrasib), anti-EGFR (cetuximab)

Parallel Pathway Inhibition: Establishing Avutometinib as the Backbone of Therapy for RAS/MAPK Pathway-Driven Tumors



Current Challenges

- Blocking RAS pathway can be circumvented through parallel pathways
 - e.g. PI3K/AKT/mTOR, FAK, RhoA, YAP
- Combinations of MEKi + AKTi have shown poor tolerability

Solutions offered with Avutometinib

- Good tolerability and early clinical data with twice weekly avutometinib opens up intermittent dosing options for combinations
- Compelling preclinical synergy data with avutometinib in combination with several key anti-cancer agents (e.g. FAKi, mTORi)
- RP2D established for avutometinib + FAKi (defactinib) and for avutometinib + mTORi (everolimus) with twice weekly regimen

Preclinical Synergy of Avutometinib in Combination with Promising Agents for Clinical Investigation

Vertical RAS/MAPK Pathway Inhibition





Parallel Pathway Inhibition

Avutometinib + pan-HERi (afatinib)

Avutometinib + SHP2i (RMC-4550)

Avutometinib + SOS1i (BI-3406)

Avutometinib + CDK4/6i (palbociclib)








Avutometinib + G12Ci (sotorasib)


Avutometinib + G12Ci (adagrasib)

Avutometinib + G12Di (MRTX1133)

Avutometinib + mTORi (everolimus)




Avutometinib Combinations in Non-Small Cell Lung Cancer

The background of the slide features a large blue area on the left. On the right side, there are several diagonal stripes in blue, teal, and orange. A horizontal teal stripe runs across the bottom of the slide.

High Unmet Need in Refractory KRAS & BRAF mt NSCLC Adenocarcinoma

NSCLC Adenocarcinoma³



KRAS Mutations Represent 25% of Lung Cancer Adenocarcinoma & BRAF Mutations Represent ~4% (EGFR 17%, ALK 7%)^{4,6}

References:

¹ Globocan, 2020

² <https://www.ncbi.nlm.nih.gov/books/NBK519578/>

³ TCGA PanCancer Atlas (cBioPortal analysis)

⁴ www.thelancet.com Vol 389 January 21, 2017


⁵ Adapted from NCCN Non-small cell lung cancer guidelines Version 3.2020

⁶ Clinical Cancer Research DOI 10.1158/1078-0432.CCR-18-2062

⁷ 50% of NSCLC is adenocarcinoma (Pakkala and Ramalingam JCI Insight 2018)

⁸ Cancer Statistics 2020, Siegel et. al. CA Cancer J Clin 2020;70:7-30

Advanced or Metastatic NSCL Cancer Recommend Histologic and Molecular Subtyping⁵



Verastem Clinical Trials:

- RAMP 203: Avutometinib + sotorasib in KRAS G12C NSCLC
- RAMP 204: Avutometinib + adagrasib in KRAS G12C NSCLC
- RAMP 202: Avutometinib + defactinib in BRAFV600E and non-V600E NSCLC


Preclinical Synergy of Avutometinib + G12C Inhibitors in KRAS G12C Models

Synergy of avutometinib + G12C inhibitors across G12C mutant NSCLC, CRC & Pancreatic cancer cell lines



Cell line	Indication	Sensitivity to G12C inhibitors	Combined Synergy Score	
			Avutometinib + sotorasib	Avutometinib + adagrasib
H2122	NSCLC	Moderately sensitive	44.7	44.6
H1373	NSCLC	Sensitive	10.0	3.4
SW1573	NSCLC	Insensitive	8.6	12.0
H358	NSCLC	Sensitive	6.9	5.4
H2030	NSCLC	Moderately sensitive	5.1	ND
SW837	CRC	Sensitive	16.1	18.5
MIAPACA2	Panc	Sensitive	2.3	5.3

ND: not determined


Avutometinib + sotorasib yields deeper and more sustained inhibition of ERK signaling pathway



Avutometinib & FAKi potentiate sotorasib efficacy in KRAS G12C NSCLC in vivo; Tumor regression in all mice with triple combination



RAMP 203: Phase I/2 Trial of Avutometinib + LUMAKRAS™ (Sotorasib) in KRAS G12C Advanced NSCLC



*may include patients with or without prior G12C therapy

NCT05074810

Abbreviations: DLT = dose-limiting toxicity; n = number of patients;
ORR = overall response rate; RP2D = recommended phase 2 dose

RAMP 204: Phase I/2 Trial of Avutometinib + KRAZATI™ (Adagrasib) in KRAS G12C Advanced NSCLC

- Patients must have a **KRAS G12C** mutation determined using validated test
- Treatment with at least 1 but no more than 3 prior systemic regimens, for Stage 3B-C or 4 NSCLC
- Patient must have received prior therapy with a **KRAS G12C** inhibitor and experience progressive disease
- **Measurable** disease according to RECIST 1.1
- **ECOG** performance status ≤ 1

Part A: Dose Evaluation
(3+3 DLT Assessment)

Part B: Dose Expansion
(Primary endpoint ORR)

Avutometinib +
Adagrasib
Dose Finding Cohorts
(n=3-6)

RP2D
Selection


Stage 1: 19 patients
(including Part A
patients) treated with
RP2D

**Stage 2: expand to 55
patients**

NCT05375994

Abbreviations: DLT = dose-limiting toxicity; n = number of patients;
ORR = overall response rate; RP2D = recommended phase 2 dose



RAMP 202: Phase 2 Trial of Avutometinib + Defactinib in BRAF mt NSCLC






Additional Avutometinib Combinations for Pancreatic, Colorectal and Melanoma

Preclinical Synergy of Avutometinib/FAK Inhibition + Chemotherapy in a KRAS/p53 pancreatic cancer mouse model



- ✓ The combination of avutometinib + FAKi induces tumor growth inhibition and increases survival but does not induce tumor regression
- ✓ Addition of chemo (gemcitabine + paclitaxel) to avutometinib/FAKi induces tumor regression

RAMP 205: Phase I/2 Trial of Avutometinib/Defactinib + GEMZAR™ (Gemcitabine)/ABRAXANE™ (Nab-paclitaxel) in Front Line Metastatic Pancreatic Cancer




NCT05669482

Abbreviations: DLT = dose-limiting toxicity; n = number of patients;
ORR = overall response rate; RP2D = recommended phase 2 dose

Combination of Avutometinib with anti-EGFR mAb Induces Tumor Regression in a KRAS mt Colorectal PDX Model

KRAS^{G12V} CRC PDX




- Avutometinib + anti-EGFR (panitumumab) induces tumor regression in a KRAS mutant CRC patient-derived xenograft model
- G12Ci + anti-EGFR (sotorasib + panitumumab and adagrasib + cetuximab) have shown partial responses in KRAS G12C CRC (Fakih et al. ESMO 2021; Weiss et al. ESMO 2021)
- **These data support the ongoing clinical evaluation of avutometinib + cetuximab (anti-EGFR) for treatment of KRAS mt CRC (NCT05200442)**

Collaboration with Marwan Fakih, City of Hope

Pachter, RAS Development Summit, 2021

Combination of Avutometinib + FAK Inhibition with Checkpoint Inhibitor Induces Tumor Regression in an IO-resistant BRAF V600E melanoma model



- Avutometinib + IO (anti-PD-1 + anti-CTLA-4) induces tumor regression in an IO-resistant syngeneic BRAF V600E melanoma model (YUMM 1.7)
- FAK inhibition deepens and sustains avutometinib-induced tumor regression
- **These data support the imminent clinical evaluation of avutometinib + pembrolizumab (anti-PD-1) for treatment of BRAF V600E melanoma**

Avutometinib Development in Multiple Combinations Across RAS Pathway-Driven Tumors with Potential Early Read-Outs in 2H 2023

Indication	Study
KRAS G12C NSCLC	RAMP 203:Avutometinib/Sotorasib combo
KRAS G12C NSCLC	RAMP 204:Avutometinib/Adagrasib combo
Pancreatic	RAMP 205:Avutometinib/Gem/Abraxane/Defactinib combo
KRAS mt NSCLC	Avutometinib/Everolimus combo*
KRAS mt CRC	Avutometinib/Cetuximab combo*
ER+ Breast	Avutometinib/Abemaciclib/fulvestrant combo*
RAS/RAF/NFI Gynecological	Avutometinib/Defactinib combo*
BRAFV600E Melanoma	Avutometinib/Pembrolizumab combo*

*Investigator Sponsored Trials

Experienced Senior Management Team



Brian Stuglik
Chief Executive Officer

- Global VP & Chief Marketing Officer – Lilly Oncology
- Founding Member – Proventus Health Solutions



Daniel Paterson
President and Chief Operating Officer

- CEO – The DNA Repair Co. (now On-Q-ity)
- PharMetrics (now IMS), Axion



Cathy Carew
Chief Organizational Effectiveness Officer

- Principal – HR Collaborative
- Ironwood, ActiveBiotics, Dynogen, Tufts Health Plan



Jonathan Pachter, Ph.D.
Chief Scientific Officer

- Head of Cancer Biology – OSI (now Astellas)
- Schering-Plough



Louis Denis, M.D.
Chief Medical Officer

- CMO, Asana BioSciences
- Boehringer-Ingelheim, Pfizer



Hagop Youssoufian, MSc, M.D.
Head of Medical Strategy

- CMO, BIND Therapeutics, EVP, Progenics,
- CMO & EVP, Ziopharm Oncology, SVP, Imclone



THANK YOU